

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSSPTA1644PNH

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page for STN Seminar Schedule - N. America  
NEWS 2 AUG 10 Time limit for inactive STN sessions doubles to 40  
minutes  
NEWS 3 AUG 18 COMPENDEX indexing changed for the Corporate Source  
(CS) field  
NEWS 4 AUG 24 ENCOMPLIT/ENCOMPLIT2 reloaded and enhanced  
NEWS 5 AUG 24 CA/CAPplus enhanced with legal status information for  
U.S. patents  
NEWS 6 SEP 09 50 Millionth Unique Chemical Substance Recorded in  
CAS REGISTRY  
NEWS 7 SEP 11 WPIDS, WPINDEX, and WPIX now include Japanese FTERM  
thesaurus  
NEWS 8 OCT 21 Derwent World Patents Index Coverage of Indian and  
Taiwanese Content Expanded  
NEWS 9 OCT 21 Derwent World Patents Index enhanced with human  
translated claims for Chinese Applications and  
Utility Models  
NEWS 10 NOV 23 Addition of SCAN format to selected STN databases  
NEWS 11 NOV 23 Annual Reload of IFI Databases  
NEWS 12 DEC 01 FRFULL Content and Search Enhancements  
NEWS 13 DEC 01 DGENE, USGENE, and PCTGEN: new percent identity  
feature for sorting BLAST answer sets  
NEWS 14 DEC 02 Derwent World Patent Index: Japanese FI-TERM  
thesaurus added  
NEWS 15 DEC 02 PCTGEN enhanced with patent family and legal status  
display data from INPADOCDB  
NEWS 16 DEC 02 USGENE: Enhanced coverage of bibliographic and  
sequence information

NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,  
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN customer  
agreement. This agreement limits use to scientific research. Use  
for software development or design, implementation of commercial  
gateways, or use of CAS and STN data in the building of commercial  
products is prohibited and may result in loss of user privileges  
and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 11:32:28 ON 11 DEC 2009

=> file medline embase biosis scisearch caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.22	0.22

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 11:32:51 ON 11 DEC 2009

FILE 'EMBASE' ENTERED AT 11:32:51 ON 11 DEC 2009

Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 11:32:51 ON 11 DEC 2009

Copyright (c) 2009 The Thomson Corporation

FILE 'SCISEARCH' ENTERED AT 11:32:51 ON 11 DEC 2009

Copyright (c) 2009 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 11:32:51 ON 11 DEC 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

=> s drug conjugat?

L1 11194 DRUG CONJUGAT?

=> s l1 and antibod?

L2 3131 L1 AND ANTIBOD?

=> s l2 and fully reducing

L3 0 L2 AND FULLY REDUCING

=> s l2 and reducing agent

L4 19 L2 AND REDUCING AGENT

=> s l4 and reoxidizing

L5 0 L4 AND REOXIDIZING

=> s l4 and oxidizing agent

L6 3 L4 AND OXIDIZING AGENT

=> dup remove l6

PROCESSING COMPLETED FOR L6

L7 3 DUP REMOVE L6 (0 DUPLICATES REMOVED)

=> s l7 and pd<20040302

1 FILES SEARCHED...

4 FILES SEARCHED...

L8 0 L7 AND PD<20040302

=> d l7 1-3 cbib abs

L7 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN

2009:487928 Document No. 150:4705290 Cysteine engineered humanized

anti-TENB2 **antibodies** and **antibody drug**

**conjugates** for cancer diagnosis and therapy. Mao, Weiguang;

Junutula, Jagath Reddy; Polakis, Paul (Genentech, Inc., USA). PCT Int.

Appl. WO 2009052249 A1 20090423, 114pp. DESIGNATED STATES: W: AE, AG,

AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO,

CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM,

GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC,

LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, MT, NE, NL, NO, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2008-US80102 20081016. PRIORITY: US 2007-981411P 20071019.

AB Cysteine engineered anti-TENB2 **antibodies** are engineered by replacing one or more amino acids of a parent anti-TENB2 **antibody** with non cross-linked, reactive cysteine amino acids. Methods of design, preparation, screening, and selection of the cysteine engineered anti-TENB2 **antibodies** are provided. Cysteine engineered anti-TENB2 **antibodies** (Ab) are conjugated with one or more drug moieties (D) through a linker (L) to form cysteine engineered anti-TENB2 **antibody-drug conjugates** having Formula I:  
Ab-(L-D)<sub>p</sub> I where p is 1 to 4. Diagnostic and therapeutic uses for cysteine engineered **antibody** drug compds. and compns. are disclosed.

L7 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN

2008:1398386 Document No. 149:575039 Cysteine-engineered humanized or chimeric anti-human MUC16 **antibodies** and **antibody drug conjugates** for diagnosis and treatment of cancer.  
Junutula, Jagath R.; Mallet, William (Genentech, Inc., USA). PCT Int. Appl. WO 2008141044 A2 20081120, 131pp. DESIGNATED STATES: W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, MT, NE, NL, NO, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2008-US62903 20080507. PRIORITY: US 2007-916657P 20070508.

AB Cysteine engineered anti-MUC16 **antibodies** are engineered by replacing one or more amino acids of a parent anti-MUC16 **antibody** with non cross-linked, reactive cysteine amino acids. Methods of design, preparation, screening, and selection of the cysteine engineered anti-MUC16 **antibodies** are provided. Cysteine engineered anti-MUC16 **antibodies** (Ab) are conjugated with one or more drug moieties (D) through a linker (L) to form cysteine engineered anti-MUC16 **antibody-drug conjugates** having Formula I:  
Ab-(L-D)<sub>p</sub> where p is 1 to 4. Diagnostic and therapeutic uses for cysteine engineered **antibody** drug compds. and compns. are disclosed.

L7 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN

2007:1390416 Document No. 148:31939 Anti-CD22 **antibodies** and immunoconjugates for diagnosis and treatment of cancer or B cell proliferative disease. Ebens, Allen J., Jr.; Gray, Alane M.; Liang, Wei-Ching; Wu, Yan; Yu, Shang-Fan (Genentech, Inc., USA). PCT Int. Appl. WO 2007140371 A2 20071206, 308 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, MT, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2007-US69889 20070529. PRIORITY: US 2006-809328P 20060530; US 2007-908941P 20070329; US 2007-911829P 20070413.

AB Anti-CD22 **antibodies** and immunoconjugates thereof are provided. Methods of using anti-CD22 **antibodies** and immunoconjugates

thereof are provided.

```
=> s conjugate
L9      331860 CONJUGATE

=> s l9 and antibod?
L10     82429 L9 AND ANTIBOD?

=> s l10 and drug
L11     22416 L10 AND DRUG

=> s l11 and reducing agent
L12     74 L11 AND REDUCING AGENT

=> s l12 and DTT
L13     8 L12 AND DTT

=> s l13 and oxidizing agent
L14     0 L13 AND OXIDIZING AGENT

=> s l13 and DTNB
L15     0 L13 AND DTNB

=> s l11 and DTT
L16     35 L11 AND DTT

=> s l16 and DTNB
L17     0 L16 AND DTNB

=> s l16 and cytotoxic agent
L18     0 L16 AND CYTOTOXIC AGENT

=> dup remove l16
PROCESSING COMPLETED FOR L16
L19     22 DUP REMOVE L16 (13 DUPLICATES REMOVED)

=> s l19 and cooling
L20     0 L19 AND COOLING

=> s selective conjugation
L21     151 SELECTIVE CONJUGATION

=> s l21 and fully reducing antibod?
L22     0 L21 AND FULLY REDUCING ANTIBOD?

=> s l21 and reducing agent
L23     0 L21 AND REDUCING AGENT

=> s (alley s?/au or trgov m?/au or sun m?/au)
L24     13588 (ALLEY S?/AU OR TORGOV M?/AU OR SUN M?/AU)

=> s l24 and conjugate
L25     152 L24 AND CONJUGATE

=> s l25 and DTT
L26     5 L25 AND DTT

=> dup remove l26
PROCESSING COMPLETED FOR L26
L27     1 DUP REMOVE L26 (4 DUPLICATES REMOVED)
```

=> d 127 cbib abs

L27 ANSWER 1 OF 1 MEDLINE on STN DUPLICATE 1  
2005502081. PubMed ID: 16173809. Reduction-alkylation strategies for the modification of specific monoclonal antibody disulfides. **Sun Michael M C**; Beam Kevin S; Cervený Charles G; Hamblett Kevin J; Blackmore Richard S; **Torgov Michael Y**; Handley Felicia G M; Ihle Nathan C; Senter Peter D; **Alley Stephen C**. (Seattle Genetics, Inc., 21823 30th Drive SE, Bothell, Washington 98021, USA. ) Bioconjugate chemistry, (2005 Sep-Oct) Vol. 16, No. 5, pp. 1282-90. Journal code: 9010319. ISSN: 1043-1802.  
Report No.: NLM-NIHMS63637; NLM-PMC2539111. Pub. country: United States. Language: English.

AB Site-specific conjugation of small molecules and enzymes to monoclonal antibodies has broad utility in the formation of **conjugates** for therapeutic, diagnostic, or structural applications. Precise control over the location of conjugation would yield highly homogeneous materials that could have improved biological properties. We describe for the first time chemical reduction and oxidation methods that lead to preferential cleavage of particular monoclonal antibody interchain disulfides using the anti-CD30 IgG1 monoclonal antibody cAC10. Alkylation of the resulting cAC10 cysteine thiols with the potent antimitotic agent monomethyl auristatin E (MMAE) enabled the assignment of drug conjugation location by purification with hydrophobic interaction chromatography followed by analysis using reversed-phase HPLC and capillary electrophoresis. These analytical methods demonstrated that treating cAC10 with reducing agents such as **DTT** caused preferential reduction of heavy-light chain disulfides, while reoxidation of fully reduced cAC10 interchain disulfides caused preferential reformation of heavy-light chain disulfides. Following MMAE conjugation, the resulting **conjugates** had isomeric homogeneity as high as 60-90%, allowing for control of the distribution of molecular species. The resulting **conjugates** are highly active both in vitro and in vivo and are well tolerated at efficacious doses.

=> s 125 and oxidizing agent

L28 1 L25 AND OXIDIZING AGENT

=> d 128 cbib abs

L28 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2009 ACS on STN  
2005:1001873 Document No. 143:304644 Proteins or antibodies conjugated with label or drug for diagnosis and treatment of cancer, immune or autoimmune disease and infection. **Alley, Stephen Charles; Torgov, Michael; Sun, Michael** (Seattle Genetics, Inc., USA). PCT Int. Appl. WO 2005084390 A2 20050915, 98 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2005-US7239 20050302. PRIORITY: US 2004-549476P 20040302.

AB A protein containing one or more activatable groups, e.g., an antibody, is subjected to partial or complete reduction of one or more such bonds to form reactive groups; the resulting protein is reacted with a drug which is reactive with some of the reactive groups, such as certain radio-metals, chelating agents, and toxins, so as to form a **conjugate** useful in, e.g., in vitro diagnosis, in vivo imaging, and therapy.

=> s 125 and fully reducing  
L29 0 L25 AND FULLY REDUCING

=> s 125 and partially reducing  
L30 0 L25 AND PARTIALLY REDUCING

=> s 125 and reducing  
L31 7 L25 AND REDUCING

=> s 131 and oxidizing  
L32 1 L31 AND OXIDIZING

=> d 132 cbib abs

L32 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2009 ACS on STN  
2005:1001873 Document No. 143:304644 Proteins or antibodies conjugated with  
label or drug for diagnosis and treatment of cancer, immune or autoimmune  
disease and infection. **Alley, Stephen Charles; Torgov,  
Michael; Sun, Michael** (Seattle Genetics, Inc., USA). PCT  
Int. Appl. WO 2005084390 A2 20050915, 98 pp. DESIGNATED STATES: W: AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,  
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,  
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,  
SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE,  
DK, ES, FI, FR, GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, NE, NL, PT, SE,  
SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2005-US7239  
20050302. PRIORITY: US 2004-549476P 20040302.

AB A protein containing one or more activatable groups, e.g., an antibody, is  
subjected to partial or complete reduction of one or more such bonds to form  
reactive groups; the resulting protein is reacted with a drug which is  
reactive with some of the reactive groups, such as certain radio-metals,  
chelating agents, and toxins, so as to form a **conjugate** useful  
in, e.g., in vitro diagnosis, in vivo imaging, and therapy.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	134.03	134.25
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-4.10	-4.10

STN INTERNATIONAL LOGOFF AT 11:41:37 ON 11 DEC 2009